

AD\_\_\_\_\_

Award Number: W81XWH-06-1-0728

TITLE: Flt-1 Function and Signaling in Breast Cancer

PRINCIPAL INVESTIGATOR: Charlotte M. Harwood

CONTRACTING ORGANIZATION: University of Massachusetts Medical Center  
Worcester, MA 01655

REPORT DATE: October 2007

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 01-10-2007		2. REPORT TYPE Annual Summary		3. DATES COVERED (From - To) 22 SEP 2006 - 21 SEP 2007	
4. TITLE AND SUBTITLE  Flt-1 Function and Signaling in Breast Cancer				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-06-1-0728	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Charlotte M. Harwood  E-Mail: Charlotte.Moriarty@umassmed.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Massachusetts Medical Center Worcester MA 01655				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The VEGF receptor Flt-1 mediates VEGF survival signaling in cancer cells and has been correlated with a high risk metastasis and relapse in breast cancer. However, the function of Flt-1 in breast cancer is not yet known. Here we report an increase in Flt-1 expression in breast tumor cells exposed to a hypoxic environment. Interestingly, no increase in Flt-1 expression was observed in pre-malignant breast epithelial cells. Furthermore, Flt-1 was identified as a target of miR-10b, which is deregulated in breast cancer. We find that expression of mature miR-10b in breast cancer cells results in significant down-regulation of Flt-1 expression. These findings will serve as a basis for functional studies of the role of Flt-1 and miR-10b in the etiology of breast cancer.					
15. SUBJECT TERMS VEGF, Flt-1, breast cancer, miR-10b					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	6	19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
<b>Introduction.....</b>	<b>4</b>
<b>Body.....</b>	<b>4</b>
<b>Key Research Accomplishments.....</b>	<b>4</b>
<b>Reportable Outcomes.....</b>	<b>4</b>
<b>Conclusion.....</b>	<b>5</b>
<b>References.....</b>	<b>5</b>


## Introduction

Vascular endothelial growth factor (VEGF) is an essential factor in breast cancer progression that signals to cancer cells in a way that facilitates their survival<sup>1,2</sup>. The survival of these cells is dependent on the interaction between VEGF and its receptor, Flt-1<sup>2,3</sup>. Recent clinical studies have correlated high levels of Flt-1 expression in breast cancer with a high risk of metastasis and relapse<sup>4,5</sup>, indicating that the interaction between VEGF and Flt-1 promotes aggressive behavior in breast cancer cells. However, the exact function of Flt-1 in breast cancer is not yet known. My overall goal is to understand the function of Flt-1 in breast cancer, with a specific focus on the role of VEGF and Flt-1 in breast cancer growth and metastasis. Specifically, I hypothesize that Flt-1 mediates VEGF survival signals and contributes to aggressive behavior in breast cancer. The scope of this project includes understanding the function of Flt-1 in breast cancer cells during tumor growth and metastasis, identifying the signaling pathways which are stimulated by Flt-1 activation, and the regulation of Flt-1 expression. This investigation into the function and signaling pathways of Flt-1 in human breast cancer will enable us to define the contribution of Flt-1 to breast cancer progression and to determine how Flt-1 signaling contributes to enhanced cancer progression. Furthermore, this work will provide the groundwork for refining current VEGF-targeted therapies, such as Bevacizumab, to target breast tumor cells directly.


## Body

To understand the function of Flt-1 in breast cancer growth and metastasis, I have been focusing on knockdown of Flt-1 expression in the SUM-159 human breast cancer cell line using RNAi. I have had limited success with consistently repressing expression of Flt-1 in this cell line, as well as others, using RNAi directed against Flt-1. Commercially available reagents have failed to inhibit Flt-1 expression using both si- and shRNA. Flt-1 knockdown using siRNA has been reported in the literature, however, I failed to reproduce the published results in my cell lines using the same siRNAs and transfection reagents. I recently entered into collaboration with Phillip Zamore, who specializes in small RNAs. Together, we have designed an siRNA which targets the 3' untranslated region of Flt-1, as opposed to the coding sequence. Preliminary studies show that this sequence is successful in knocking down Flt-1 protein expression by western immunoblot, as compared to a non-targeting siRNA (Figure 1).

To gain further insight into Flt-1 function, I have begun to examine the impact of hypoxia on Flt-1 expression in the MCF-10A series of breast cancer cells. In all four stages, ranging from normal breast epithelial cells to metastatic breast cancer cells, Flt-1 expression was assessed by western immunoblot following exposure to a severely hypoxic environment (0.1% O<sub>2</sub>) for 24 hours. Flt-1 levels were enhanced in the tumorigenic cell lines MCF-10CA1h and MCF-10CA1a, but did not change in normal MCF-10A or pre-malignant MCF-10AT cells, as compared to control cells maintained in normoxia for the same period of time (Figure 2).




**Figure 1.** Flt-1 siRNA decreases Flt-1 protein expression in SUM-159 cells.



**Figure 2.** Expression of Flt-1 protein increases following 24 hours in a hypoxic environment (0.1% O<sub>2</sub>) in tumorigenic variants of MCF-10A cells, but not in non-tumorigenic or premalignant variants.


To address Flt-1 signaling in breast cancer, I proposed to use an Flt-1/CSF-1 chimeric receptor composed of the Flt-1 intracellular and transmembrane domains fused to the CSF-1 extra-cellular domain<sup>6</sup>. However, CSF-1R was recently found to be commonly expressed in both normal breast epithelium and breast tumors. Additionally, CSF-1R is reported to play a significant role in normal breast development, as well as in mammary tumor progression to metastasis<sup>7,8</sup>. I examined several human breast cancer cell lines using reverse-transcriptase RT-PCR and found them all to express CSF-1R and/or its ligand CSF-1 (Figure 3). As a result, this

chimeric receptor is not suitable for use in these cells, as basal stimulation by endogenous ligand would present a confounding factor. I am currently investigating other options for examining Flt-1 signaling in breast cancer.



**Figure 3.** Expression of a CSF-1 and CSF-1 receptor in human breast cancer cell lines by RT-PCR.

Finally, I have begun to examine the regulation of Flt-1 expression in breast cancer to determine what factor or factors are causing Flt-1 levels to increase in cancer cells. Recent publications have implicated miR-10b as one of several microRNAs deregulated in breast cancer<sup>9,10</sup>. Flt-1 is one of many genes predicted by computer algorithms to be a target of miR-10b, therefore I hypothesized that miR-10b regulates the expression of Flt-1 in breast carcinoma. Utilizing a miR-10b mimic and a mutant miR-10b containing a single base-pair substitution in the seed region as a control, I found that expression of miR-10b results in a downregulation of Flt-1 protein by western immunoblot in SUM-159 breast cancer cells lacking endogenous miR-10b. Expression of the miR-10b seed mutant or a non-targeting control mimic had no effect on Flt-1 levels as compared to untreated cells (Figure 4). I plan to continue these studies, using ectopic expression of miR-10b in breast cancer cells as well as antisense for miR-10b silencing, to examine the function of miR-10b in breast cancer and its regulation of Flt-1 expression. Furthermore, I will assess the regulation of Flt-1 by several other miRNAs that are predicted to target the Flt-1 3' UTR.



**Figure 4.** Expression of a miR-10b mimic results in downregulation of Flt-1 protein expression.

### Key Research Accomplishments

- Hypoxia increases expression of Flt-1 in tumorigenic breast cell lines, but not in pre-malignant breast epithelial cells
- Flt-1 expression in breast cancer is regulated, at least in part, by miR-10b
- Presentation of work at lab meetings and departmental data club
- Attendance at UMass Memorial's monthly Breast Cancer Conference
- Attendance at the AACR Special Conference: The Role of Non-Coding RNAs in Cancer

### Reportable Outcomes

None

## Conclusion

In conclusion, hypoxia was found to regulate expression of Flt-1 in breast cancer and Flt-1 was validated as a target of miR-10b, a microRNA which is deregulated in breast cancer. Future work will address the significance of these findings with specific regard to Flt-1 function, as well as to the function of miR-10b, in breast cancer growth and metastasis. The mechanisms of Flt-1 regulation and Flt-1 signaling in breast cancer will also be addressed. An understanding of how Flt-1 contributes to growth and progression in breast cancer will have broad implications for other common epithelial cancers and will provide a foundation of knowledge that will potentially enable the development of therapeutic strategies aimed at targeting this important pathway in breast cancer.

## References

1. Bachelder, R. E., Crago, A., Chung, J., Wendt, M. A., Shaw, L. M., Robinson, G., and Mercurio, A. M. Vascular endothelial growth factor is an autocrine survival factor for neuropilin-expressing breast carcinoma cells. *Cancer Res.*, 61: 5736-5740, 2001.
2. Bates, R. C., Goldsmith, J. D., Bachelder, R. E., Brown, C., Shibuya, M., Oettgen, P., and Mercurio, A. M. Flt-1-dependent survival characterizes the epithelial-mesenchymal transition of colonic organoids. *Curr.Biol.*, 13: 1721-1727, 2003.
3. Das, B., Yeger, H., Tsuchida, R., Torkin, R., Gee, M. F., Thorner, P. S., Shibuya, M., Malkin, D., and Baruchel, S. A hypoxia-driven vascular endothelial growth factor/Flt1 autocrine loop interacts with hypoxia-inducible factor-1alpha through mitogen-activated protein kinase/extracellular signal-regulated kinase 1/2 pathway in neuroblastoma. *Cancer Res.*, 65: 7267-7275, 2005.
4. 't Veer, L. J., Dai, H., van de Vijver, M. J., He, Y. D., Hart, A. A., Mao, M., Peterse, H. L., van der, K. K., Marton, M. J., Witteveen, A. T., Schreiber, G. J., Kerkhoven, R. M., Roberts, C., Linsley, P. S., Bernards, R., and Friend, S. H. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*, 415: 530-536, 2002.
5. Meunier-Carpentier, S., Dales, J. P., Djemli, A., Garcia, S., Bonnier, P., Andrac-Meyer, L., Lavaut, M. N., Allasia, C., and Charpin, C. Comparison of the prognosis indication of VEGFR-1 and VEGFR-2 and Tie2 receptor expression in breast carcinoma. *Int.J.Oncol.*, 26: 977-984, 2005.
6. Rahimi, N., Dayanir, V., and Lashkari, K. Receptor chimeras indicate that the vascular endothelial growth factor receptor-1 (VEGFR-1) modulates mitogenic activity of VEGFR-2 in endothelial cells. *J.Biol.Chem.*, 275: 16986-16992, 2000.
7. Yee LD, Liu L. The constitutive production of colony stimulating factor 1 by invasive human breast cancer cells. *Anticancer Res.* 20(6B):4379-83, 2000.
8. Lin EY, Nguyen AV, Russell RG, Pollard JW. Colony-stimulating factor 1 promotes progression of mammary tumors to malignancy. *J Exp Med.* 193(6):727-40, 2001.
9. Iorio MV, Ferracin M, Liu CG, Veronese A, Spizzo R, Sabbioni S, Magri E, Pedriali M, Fabbri M, Campiglio M, Menard S, Palazzo JP, Rosenberg A, Musiani P, Volinia S, Nenci I, Calin GA, Querzoli P, Negrini M, Croce CM. MicroRNA gene expression deregulation in human breast cancer. *Cancer Res.* 65(16):7065-70, 2005.
10. Ma L, Teruya-Feldstein J, Weinberg RA. Tumour invasion and metastasis initiated by microRNA-10b in breast cancer. *Nature.* 449(7163):682-8, 2007.